

COMMUTING ELEMENTS, SIMPLICIAL SPACES AND FILTRATIONS OF CLASSIFYING SPACES

A. ADEM*, F. R. COHEN**, AND E. TORRES GIESE

ABSTRACT. Let Γ^q denote the q -th stage of the descending central series of the free group on n generators F_n . For each $q \geq 2$ and every topological group G , a simplicial space $B_*(q, G)$ is constructed where $B_n(q, G) = \text{Hom}(F_n/\Gamma^q, G)$ and the realizations $B(q, G) = |B_*(q, G)|$ filter the classifying space BG . In particular for $q = 2$ this yields a single space $B(2, G)$ assembled from all the n -tuples of commuting elements in G . Homotopy properties of the $B(q, G)$ are considered for finite groups, including their description as homotopy colimits. Cohomology calculations are provided for compact Lie groups. The spaces $B(2, G)$ are described in detail for transitively commutative groups. Stable homotopy decompositions of the $B(q, G)$ are also provided with a formula giving the cardinality of $\text{Hom}(F_n/\Gamma^q, G)$ for finite discrete groups G (and thus the cardinality of $\text{Hom}(\mathbb{Z}^n, G)$ in case $q = 2$) in terms of the ranks of the homology groups for the associated filtration quotients of $B(q, G)$. Specific calculations for $H_1(B(q, G); \mathbb{Z})$ are shown to be delicate in case G is finite of odd order in the sense that resulting topological properties (which are not yet fully understood) are equivalent to the Feit–Thompson theorem.

CONTENTS

1. Introduction	2
2. On the Simplicial Structure of $\text{Hom}(F_n/\Gamma^r, G)$	4
3. $B(q, G)$ as a Colimit and Homotopy Colimit	13
4. The Homology of $B(q, G)$ when G is Finite	17
5. The Spaces $B(q, G)$ for Compact, Connected Lie Groups	19
6. $B(2, G)$ for Transitively Commutative Groups	23
7. Stable Splittings and the Cardinality of $\text{Hom}(F_n/\Gamma^q, G)$ for Finite Groups	34
References	38

Date: January 12, 2009.

*Partially supported by NSERC and NSF.

**Partially supported by DARPA and the NSF.

1. INTRODUCTION

Let G denote a topological group, its classifying space BG plays a central role in algebraic topology with important applications to bundle theory and cohomology of groups. In this paper a filtration of BG is introduced by using the descending central series of the free groups. A basic observation is that if F_n is the free group on n generators with Γ^q the q -th stage of its descending central series, then the spaces of homomorphisms $\text{Hom}(F_n/\Gamma^q, G)$ can be assembled to form simplicial spaces with realizations $B(q, G)$ which filter the usual classifying space BG . In other words there are inclusions

$$B(2, G) \subset B(3, G) \subset \cdots \subset B(q, G) \subset B(q+1, G) \subset \cdots \subset B(\infty, G) = BG$$

where each term is constructed from the simplicial spaces associated to terms in the descending central series of the free group. This naturally gives rise to a functorial construction on topological groups $G \mapsto B(q, G)$. In fact the construction provided here affords a principal G -bundle $E(q, G) \rightarrow B(q, G)$ which fits into a commutative diagram, namely there are natural morphisms of principal G -bundles

$$\begin{array}{ccccc} E(q, G) & \xrightarrow{e_q} & E(q+1, G) & \longrightarrow & EG \\ \downarrow p & & \downarrow p & & \downarrow \\ B(q, G) & \xrightarrow{b_q} & B(q+1, G) & \longrightarrow & BG \end{array}$$

and the maps e_q, b_q yield a natural filtration of subspaces for EG and BG .

In this paper the basic properties of these spaces are studied; they turn out to have many good features, and a lot of work remains to be done to fully understand them. Unlike the classical situation, the spaces $E(q, G)$ are not necessarily contractible. Thus the fibration $E(q, G) \rightarrow B(q, G) \rightarrow BG$ can be quite interesting, in particular for G a finite group it gives rise to natural representations of G on the homology of $E(q, G)$. For a connected Lie group G there is a homotopy equivalence for all $q \geq 2$:

$$G \times \Omega(E(q, G)) \simeq \Omega B(q, G).$$

In terms of cohomology the following results are proved in §2:

- (1) If G is a finite group, then for any $q \geq 2$, the map on mod- p cohomology

$$\phi^* : H^*(BG; \mathbb{F}_p) \rightarrow H^*(B(q, G); \mathbb{F}_p)$$

has a nilpotent kernel.

- (2) If G is a finite group with mod p cohomology detected by subgroups of nilpotence class less than q , then $H^*(BG; \mathbb{F}_p) \rightarrow H^*(B(q, G); \mathbb{F}_p)$ is a monomorphism.
- (3) Let G denote a compact, connected Lie group with maximal torus $T \subset G$. Assume that the spaces $Hom(\mathbb{Z}^n, G)$ are all path connected. There is an isomorphism

$$H^*(B(2, G); \mathbb{Q}) \cong H^*(G/T \times BT; \mathbb{Q})^{W(G)}$$

which is compatible with the well-known isomorphism $H^*(BG; \mathbb{Q}) \cong H^*(BT; \mathbb{Q})^{W(G)}$, where $W(G)$ denotes the Weyl group of T in G .

Let $\mathcal{N}_q(G)$ denote the family of all subgroups of G of nilpotence class less than q and denote $G(q) = \operatorname{colim}_{A \in \mathcal{N}_q(G)} A$. The sequence of groups $G(2), G(3), G(4), \dots$ captures information relevant to the subgroups of G of increasing nilpotence class; note that $G(\infty) = G$. If G is a finite group, then for any $q \geq 2$, it can be shown that

$$B(q, G) \simeq \operatorname{hocolim}_{A \in \mathcal{N}_q(G)} BA$$

and this can be used to show

Theorem 1.1. *There is a natural fibration $B(q, G) \rightarrow BG(q)$ with fiber a simply-connected finite dimensional complex K_q .*

A basic question is that of determining under what conditions the fibre K_q is contractible and $B(q, G) \simeq BG(q)$. This condition is verified for a class of groups, where an associated graph constructed from maximal subgroups of nilpotence class less than q turns out to be a tree. In particular this yields that for *transitively commutative groups* (TC groups) $B(2, G) \simeq BG(2)$, and explicit computations can be provided. In particular if G is a TC group then the fundamental group of $E(2, G)$ is a free group of rank

$$N_G = 1 - |G : Z(G)| + \left(\sum_{1 \leq i \leq k} |G : Z(G)| - |G : C_G(a_i)| \right)$$

where the centralizers $C_G(a_i)$, $i = 1, \dots, k$, represent a minimal collection of centralizers covering G (see §6 for precise definitions). Moreover it can be shown that

Proposition 1.2. *If G is a finite TC group with trivial center, then there is a stable homotopy equivalence*

$$B(2, G) \simeq \bigvee_{p || G|} \bigvee_{P \in \operatorname{Syl}_p(G)} BP.$$

For example, the alternating group A_5 is a TC group and there is a homotopy equivalence

$$B(2, A_5) \simeq [\bigvee^5 B(\mathbb{Z}/2 \times \mathbb{Z}/2)] \bigvee [\bigvee^{10} B\mathbb{Z}/3] \bigvee [\bigvee^6 B\mathbb{Z}/5].$$

In the last section stable splittings associated to the individual simplicial spaces are discussed. Let $S_k(j, q, G) \subset \text{Hom}(F_k/\Gamma^q, G) \subset G^k$ denote the subspace that consists of k -tuples which have 1 in at least j coordinates. A Lie group G is said have *cofibrantly filtered elements* if the natural inclusions $I_j : S_k(j, q, G) \rightarrow S_k(j-1, q, G)$ are cofibrations for all k, q and j for which both spaces are non-empty. Then it follows from general results about simplicial spaces (see [2]) extending those in [3] that there are stable homotopy decompositions described by¹:

Theorem 1.3. *If G has cofibrantly filtered elements, then there are homotopy equivalences*

$$\Sigma \text{Hom}(F_n/\Gamma^q, G) \rightarrow \bigvee_{1 \leq k \leq n}^{\binom{n}{k}} \Sigma \bigvee \text{Hom}(F_k/\Gamma^q, G)/S_k(q, G).$$

Moreover, the natural filtration quotients

$$E_k^0(B(q, G)) = F_k B(q, G)/F_{k-1} B(q, G)$$

of the geometric realization $B(q, G)$ are stably homotopy equivalent to the summands

$$\text{Hom}(F_k/\Gamma^q, G)/S_k(q, G).$$

Analogous constructions can be done using the p -descending central series for the free groups and using the free pro- p -groups. These variations lead to local versions of the constructions in this paper. It is also possible to construct a simplicial space using the spaces of representations $\text{Hom}(F_n/\Gamma^q, G)/G^{ad}$ (here the quotient is by the conjugation action). The resulting spaces, denoted $\overline{E}(q, G)$ and $\overline{B}(q, G)$ will have equally interesting properties and will be discussed in a sequel to this paper.

The third author would like to thank Fred Cohen for his hospitality and support given during his stay at the University of Rochester in Spring 2008.

2. ON THE SIMPLICIAL STRUCTURE OF $\text{Hom}(F_n/\Gamma^r, G)$

The goal of this section is to define a family of simplicial spaces assembled using spaces of homomorphisms. In fact, this family will yield a filtration of the bar construction of the

¹If X and Y are pointed topological spaces, then $X \vee Y$ denotes their one point union in the product $X \times Y$ and ΣX denotes the suspension of X

classifying space of a group and it will be parametrized by the lower central series of the free group.

The following is a standard definition from group theory.

Definition 2.1. Let Q be a group, define a chain of subgroups $\Gamma^r(Q)$ inductively: $\Gamma^1(Q) = Q$; $\Gamma^{i+1}(Q) = [\Gamma^i(Q), Q]$. By convention, $\Gamma^\infty(Q) = \{1\}$. The descending central series of Q is the normal series

$$\dots \subset \Gamma^{i+1}(Q) \subset \Gamma^i(Q) \subset \dots \subset \Gamma^2(Q) \subset \Gamma^1(Q) = Q.$$

A discrete group Q is said to be *nilpotent* if there is some integer m such that $\Gamma^{m+1}(Q) = \{1\}$. The least such integer m is called the *class* of Q . For example every finite p -group is nilpotent.

When the context is clear Γ^r will be used to denote these normal subgroups of Q . Let $F_n = F[e_0, e_1, \dots, e_{n-1}]$ denote the free group on n -letters. Fix an integer q and let Γ^q denote the q -stage of the descending central series for F_n , as above.

Let G denote a topological group; in this paper it will be assumed throughout that: (i) G is locally compact as well as Hausdorff; and that (ii) $1 \in G$ is a non-degenerate basepoint. Consider the set $\text{Hom}(F_n/\Gamma^q, G)$ of homomorphisms $f: F_n \rightarrow G$ which descend to homomorphisms $\bar{f}: F_n/\Gamma^q \rightarrow G$, i.e. such that $f(\Gamma^q) = 1$. An element of $\text{Hom}(F_n/\Gamma^q, G)$ is specified by functions from the set $\{e_0, e_1, \dots, e_{n-1}\}$ to G . Thus an element in $\text{Hom}(F_n/\Gamma^q, G)$ is identified as an ordered n -tuple $(x_0, x_1, \dots, x_{n-1})$ with $f(e_i) = x_i$. The space $\text{Hom}(F_n/\Gamma^q, G)$ is topologized as a subspace of G^n with the naturally inherited topology for general G .

Two simplicial spaces associated to these homomorphisms will now be introduced.

Definition 2.2. Let

$$E_n(q, G) = G \times \text{Hom}(F_n/\Gamma^q, G) \subset G^{n+1},$$

and define $d_i: E_n(q, G) \rightarrow E_{n-1}(q, G)$ for $0 \leq i \leq n$ and $s_j: E_n(q, G) \rightarrow E_{n+1}(q, G)$, for $0 \leq j \leq n$, given by

$$d_i(g_0, \dots, g_n) = \begin{cases} (g_0, \dots, g_i \cdot g_{i+1}, \dots, g_n) & 0 \leq i < n \\ (g_0, \dots, g_{n-1}) & i = n \end{cases}$$

and

$$s_j(g_0, \dots, g_n) = (g_0, \dots, g_i, e, g_{i+1}, \dots, g_n)$$

for $0 \leq j \leq n$.

Definition 2.3. Similarly, let

$$B_n(q, G) = \text{Hom}(F_n/\Gamma^q, G)$$

with maps d_i and s_j defined in the same way, except that the first coordinate g_0 is omitted and the map d_0 takes the form $d_0(g_1, \dots, g_n) = (g_2, \dots, g_n)$.

Lemma 2.4. *The maps d_i, s_j defined on the spaces $E_n(q, G)$ and $B_n(q, G)$ are well-defined and equip them with the structure of simplicial spaces.*

Proof. The maps d_i and s_i are well-defined since they are induced by group homomorphisms between free groups and from the fact that if $f : A \rightarrow B$ is a group homomorphism then $f(\Gamma^q A) \subseteq \Gamma^q B$. The simplicial identities follow (as in the classical case for the bar construction) from the definition of the homomorphisms inducing the maps d_i and s_j . $\square$

Note that the map $G^{n+1} \rightarrow G^n$ that projects the last n coordinates onto G^n defines a simplicial map

$$p : E_*(q, G) \rightarrow B_*(q, G)$$

Moreover, G acts from the left on $E_n(q, G)$ by multiplication on the first coordinate

$$g(g_0, g_1, \dots, g_n) = (gg_0, g_1, \dots, g_n)$$

and this action makes $E_*(q, G)$ into a G -simplicial space; that is, the action of G commutes with the face and degeneracy maps. This action is free and its degree-wise orbit space is homeomorphic to $B_*(q, G)$.

Recall the notion of geometric realization for a simplicial space.

Definition 2.5. The geometric realization of a simplicial space Z_* is the following topological space

$$|Z_*| := \coprod_{n \geq 0} Z_n \times \Delta^n / \sim$$

where Δ^n denotes the n -simplex and the equivalence relation $\sim$ is defined as follows. Identify $(x, \delta_i t) \in X_n \times \Delta^n$ with $(d_i x, t) \in X_{n-1} \times \Delta^{n-1}$ for any $x \in X_n$, $t \in \Delta^{n-1}$ and $(x, \sigma_j t) \in X_n \times \Delta^n$ with $(s_j x, t) \in X_{n+1} \times \Delta^{n+1}$ for any $x \in X_{n-1}$ and $t \in \Delta^{n+1}$. The topology on $|Z_*|$ is the quotient topology.

Definition 2.6. The realization $|Z_*|$ of a simplicial space has a canonical filtration defined as follows: $F_r |Z_*| \subset |Z_*|$ is the image of $\coprod_{0 \leq i \leq r} Z_i \times \Delta^i$ in the quotient space. The associated graded space is defined as $E_j^0(|Z_*|) = F_j |Z_*| / F_{j-1} |Z_*|$.

The following definition will be used later and will help the understanding of the local structure of the map p on the geometric realizations.

Definition 2.7. Let $F_j G^n$ be the subspace of G^n that consists of n -tuples with at least j coordinates equal to e , define

$$S_n(j, q, G) = \text{Hom}(F_n/\Gamma^q, G) \cap F_j G^n$$

Proposition 2.8. *The map $p : |E_*(q, G)| \rightarrow |B_*(q, G)|$, induced by the maps p_n , is a principal G -bundle.*

Proof. The relations imposed by the face and degeneracy maps show that

$$F_n |E_*(q, G)| - F_{n-1} |E_*(q, G)| = G \times (\text{Hom}(F_n/\Gamma_q, G) - S_n(1, q, G)) \times (\Delta_n - \partial\Delta_n)$$

whereas

$$F_n |B_*(q, G)| - F_{n-1} |B_*(q, G)| = (\text{Hom}(F_n/\Gamma_q, G) - S_n(1, q, G)) \times (\Delta_n - \partial\Delta_n)$$

The map p restricts to the projection between these subspaces, and so it is a G -bundle as long as the identity of G is a non-degenerate base-point. □

Remark. Notice that the spaces $E_*(q, G)$ and $B_*(q, G)$ are simplicial subspaces of the bar construction of G . In fact, $|E_*(\infty, G)| = EG$, $|B_*(\infty, G)| = BG$ and p is the standard map $EG \rightarrow BG$. In other words, this yields Milgram's model for the universal principal G -bundle $EG \rightarrow BG$ (see [12] and [4], page 49).

Proposition 2.9. *Let G be a group.*

- 1) *The natural surjection $F_n/\Gamma^{q+1} \rightarrow F_n/\Gamma^q$ induces a map of simplicial spaces compatible with the simplicial map p_n ; that is, the diagram*

$$\begin{array}{ccc} E_n(q, G) & \xrightarrow{e_q} & E_n(q+1, G) \\ \downarrow p_n & & \downarrow p_n \\ B_n(q, G) & \xrightarrow{b_q} & B_n(q+1, G) \end{array}$$

commutes.

- 2) *There are natural morphisms of principal G -bundles*

$$\begin{array}{ccccc} |E_*(q, G)| & \xrightarrow{e_q} & |E_*(q+1, G)| & \longrightarrow & EG \\ \downarrow p & & \downarrow p & & \downarrow \\ |B_*(q, G)| & \xrightarrow{b_q} & |B_*(q+1, G)| & \longrightarrow & BG \end{array}$$

- 3) The maps e_q, b_q yield a natural filtration of subspaces for EG and BG .
 4) If G is a nilpotent group of class c , then $|E_*(c+1, G)| = EG$, $|B_*(c+1, G)| = BG$ and the filtration is finite. If G is finitely generated, then the filtration is finite if and only if G is nilpotent.

Proof. Part (1) and (2) are immediate since the maps e_q, b_q are induced by group homomorphisms, whereas part (3) follows from (2). If G is nilpotent of class c , then $\Gamma^{c+1}(G) = \{1\}$ and thus $\text{Hom}(F_n/\Gamma^{c+1}, G) = G^n$ for all n . Hence $|E_*(c+1)| = EG$ and $|B_*(c+1)| = BG$. If $|B_*(q, G)| = BG$ for some q , then for all n

$$F_n|B_*(q, G)| - F_{n-1}|B_*(q, G)| = F_n|B_*(\infty, G)| - F_{n-1}|B_*(\infty, G)|$$

Suppose that G is generated by m elements and that $\phi : F_m \rightarrow G$ is a homomorphism onto the generators of G . Furthermore,

$$\text{Hom}(F_m/\Gamma_q, G) - S_m(1, q, G) = \text{Hom}(F_m, G) - S_m(1, \infty, G)$$

and $\phi \in \text{Hom}(F_m, G) - S_m(1, \infty, G)$. Therefore $\Gamma^q(G) = \phi(\Gamma^q) = \{1\}$ and G is nilpotent. $\square$

Definition 2.10. The geometric realizations of $E_*(q, G)$ and $B_*(q, G)$ will be denoted by $E(q, G)$ and $B(q, G)$ respectively.

The map $B(q, G) \rightarrow BG$ can be identified with the classifying map associated to the principal G -bundle $E(q, G) \rightarrow B(q, G)$, whence there is a fibration

$$E(q, G) \rightarrow B(q, G) \rightarrow BG.$$

The functor from topological groups to topological spaces, given by

$$G \mapsto B(q, G)$$

has a number of interesting features, some of them analogous to the classifying space functor but there are some important differences. Given a homomorphism of topological groups $f : G \rightarrow H$ the notation

$$E(q, f) : E(q, G) \rightarrow E(q, H) \text{ and } B(q, f) : B(q, G) \rightarrow B(q, H)$$

will be used to denote the induced maps and $p(q, G) : E(q, G) \rightarrow B(q, G)$ will denote the natural map from now on.

The following proposition is a list of some of their basic properties.

Proposition 2.11. *The functor $B(q, G)$ satisfies the following properties:*

- (1) If $q \geq 2$ and H is a topological group of nilpotency class less than q , then $B(q, H) = BH$, the usual classifying space.

- (2) If G is a finite group then there exists an N that depends on G such that $B(q, G) = B(N, G)$ for all $q \geq N$.
- (3) If $q \geq 2$ and $\iota: H \rightarrow G$ is a homomorphism where H is a group of nilpotency class less than q , then there is a commutative diagram

$$\begin{array}{ccc} BH & \xrightarrow{1} & BH \\ \downarrow & & \downarrow B(\iota) \\ B(q, G) & \longrightarrow & BG. \end{array}$$

Thus if BG is a stable retract of BH , then BG is a stable retract of $B(q, G)$.

- (4) If $q \geq 2$ and Z is a central subgroup of G , then the natural action map

$$\mu: Z \times G \rightarrow G$$

induces an action

$$\mu: BZ \times B(q, G) \rightarrow B(q, G)$$

together with a commutative diagram

$$\begin{array}{ccc} BZ \times BZ & \xrightarrow{1} & BZ \\ \downarrow & & \downarrow I \\ BZ \times B(q, G) & \xrightarrow{\mu} & B(q, G). \end{array}$$

- (5) If π is finitely generated, the functor $\text{Hom}(\pi, G)$ commutes with colimits in G , i.e.

$$\text{Hom}(\pi, \varinjlim_{\alpha} G_{\alpha}) \rightarrow \varinjlim_{\alpha} \text{Hom}(\pi, G_{\alpha})$$

is a homeomorphism. Therefore there are homeomorphisms

$$B(q, \varinjlim_{\alpha} G_{\alpha}) \cong \varinjlim_{\alpha} B(q, G_{\alpha}).$$

Proof. These properties are easy to verify, in particular the stable value $B(N, G)$ for the spaces $B(q, G)$ can be explained as follows. Let N denote the minimal integer such that the class of nilpotence for any nilpotent subgroup of G is strictly less than N . It suffices to show that

$$\text{Hom}(F_n/\Gamma^N, G) = \text{Hom}(F_n/\Gamma^{N+t}, G)$$

for any $t \geq 1$. Given a homomorphism $\phi: F_n \rightarrow G$, where F_n is a free group of rank n , let $H = \text{Im}\phi$. Then ϕ will descend to a map $F_n/\Gamma^{N+t} \rightarrow G$ if and only if the nilpotence class of H is less than $N + t$. However, given that H is nilpotent, its nilpotence class is in fact less

than N ; hence ϕ represents an element in $\text{Hom}(F_n/\Gamma^N, G)$ and the proof is complete. Note that if G is nilpotent then the stable value is BG . $\square$

Recall that the map $c_g : BG \rightarrow BG$ induced by conjugation by $g \in G$ is homotopic to the identity. It turns out that for $B(q, G)$ this need not be the case. The reason is that $B_*(q, G)$ splits into a lattice of nerves of categories so that conjugation may permute this lattice. On the other hand, this difference indicates that when comparing the cohomology of BG with that of $B(q, G)$ it is only necessary to look at the invariant part of the cohomology of $B(q, G)$ as the following diagram

$$\begin{array}{ccc} B(q, G) & \longrightarrow & BG \\ \downarrow c_g & & \downarrow c_g \\ B(q, G) & \longrightarrow & BG \end{array}$$

commutes. Hence the map $H^*(BG; \mathbb{Z}) \rightarrow H^*(B(q, G); \mathbb{Z})$ factors through $H^*(B(q, G); \mathbb{Z})^G$.

The next proposition deals with basic facts about the cohomology of the spaces $B(q, G)$. These have close connections to properties of group cohomology.

Proposition 2.12. *The functors $B(q, G)$ satisfy the following properties:*

- (1) *If G is a finite group with mod p cohomology detected by subgroups of nilpotence class less than q , then $H^*(BG; \mathbb{F}_p) \rightarrow H^*(B(q, G); \mathbb{F}_p)$ is a monomorphism.*
- (2) *If G is a finite group, then for any $q \geq 2$, the map on mod- p cohomology*

$$\phi^* : H^*(BG; \mathbb{F}_p) \rightarrow H^*(B(q, G); \mathbb{F}_p)$$

is a monomorphism modulo nilpotent elements.

- (3) *The spectral sequence obtained from the skeletal filtration satisfies*

$$E_{s,t}^1 = \oplus H_{s+t}(F_s B(q, G), F_{s-1} B(q, G); \mathbb{Z})$$

and abuts to $H_{s+t}(B(q, G); \mathbb{Z})$. In case $q = \infty$, and coefficients are in a field $\mathbb{F}$, this spectral sequence is the bar spectral sequence with

$$E_{s,t}^2 = \text{Tor}_{s,t}^{H^*(G)}(\mathbb{F}, \mathbb{F})$$

which abuts to $H_{s+t}(BG; \mathbb{F})$.

Proof. The first statement follows immediately from parts (1) and (2) of the previous proposition. For the second statement it suffices to prove it for $q = 2$. Recall the result due to Quillen and Venkov, namely that if $A_p(G)$ denotes the poset of non-trivial elementary

abelian p -subgroups of G with morphisms induced by inclusion and conjugation, then the map induced by the restrictions

$$Q : H^*(BG; \mathbb{F}_p) \rightarrow \lim_{E \in A_p(G)} H^*(E; \mathbb{F}_p)$$

has nilpotent kernel (see [17] and [4], page 146). Note, however, that this factors through the invariants of the mod p cohomology of $B(q, G)$ for any $q \geq 2$; hence any element in the kernel of $\phi^* : H^*(BG; \mathbb{F}_p) \rightarrow H^*(B(2, G); \mathbb{F}_p)$ must be nilpotent, as it is necessarily in the kernel of the map Q .

The statement about the spectral sequence is routine and the identification with the bar spectral sequence in the classical case is well-known (see [15]). $\square$

Instead of considering the descending central series of a group, it is also possible to consider its p -descending central series, defined as follows:

Definition 2.13. Let p denote a prime number and Q a group; define a chain of subgroups $\Gamma_p^r(Q)$ inductively: $\Gamma_p^1(Q) = Q$; $\Gamma_p^{i+1}(Q) = [\Gamma_p^i(Q), Q](\Gamma_p^i(Q))^p$. By convention, $\Gamma_p^\infty(Q) = \{1\}$. The p -descending central series of Q is the normal series

$$\cdots \subset \Gamma_p^{i+1}(Q) \subset \Gamma_p^i(Q) \subset \cdots \subset \Gamma_p^2(Q) \subset \Gamma_p^1(Q) = Q.$$

A discrete group Q is said to be p -nilpotent if there is some integer m such that $\Gamma_p^{m+1}(Q) = \{1\}$. The least such integer is called the p -class of Q . Given the free group F_n on n letters, denote by Γ_p^q the q -th stage of the descending central series for F_n .

There are some important differences here, in particular $\Gamma_p^i/\Gamma_p^{i+1}$ is an elementary abelian p -subgroup and the quotient groups F_n/Γ_p^i are all finite p -groups. Just as before, they can be assembled to yield simplicial spaces $E_*(q, G, p)$ and $B_*(q, G, p)$ where G is a topological group. The basic fact used is that if $f : A \rightarrow B$ is a group homomorphism then $f(\Gamma_p^i(A)) \subset \Gamma_p^i(B)$. We can take their geometric realization and thus obtain spaces $E(q, G, p)$ and $B(q, G, p)$; they will now exhibit properties specific to the prime p . For example $B(2, G, p)$ is a space assembled from the p -elementary abelian subgroups in G . Here is an example of how these spaces capture p -local information.

Proposition 2.14. *Let G denote a finite group such that its mod p cohomology is detected on p -elementary abelian subgroups; then the map $B(2, G, p) \rightarrow BG$ induces an injective map in mod p cohomology.*

The proof follows from restricting to elementary abelian subgroups. Note the inclusions

$$B(2, G, p) \subset B(2, G) \subset BG$$

indeed detecting on $B(2, G, p)$ is much sharper than doing it with $B(2, G)$.

Example 2.15. Let $G = \Sigma_n$, the symmetric group on n letters. Then it is well known (see [4], Ch. VII) that its mod 2 cohomology is detected on elementary abelian 2-groups. Thus it follows that for all $n \geq 1$

$$H^*(\Sigma_n; \mathbb{F}_2) \hookrightarrow H^*(B(2, \Sigma_n, 2); \mathbb{F}_2)$$

is a monomorphism.

The group G acts by simultaneous conjugation on the coordinates of the spaces $E_n(q, G)$ and $B_n(q, G)$. It can be easily verified that the face and degeneracy maps are all equivariant with respect to this *adjoint* action, as are the projection maps $p_n : E_n(q, G) \rightarrow B_n(q, G)$. Hence the quotients will form simplicial spaces:

Definition 2.16.

$$\overline{E}_*(q, G) = E_*(q, G)/G^{ad} \quad \overline{B}_*(q, G) = B_*(q, G)/G^{ad}$$

with realizations

$$\overline{E}(q, G) = |\overline{E}_*(q, G)| \quad \overline{B}(q, G) = |\overline{B}_*(q, G)|$$

In particular the space $\overline{B}(q, G)$ can be thought of as the realization of a simplicial space assembled from spaces of *representations* $\text{Rep}(F_n/\Gamma^q, G) = \text{Hom}(F_n/\Gamma^q, G)/G$. The action of G by conjugation on the simplicial spaces gives rise to actions induced by conjugation on the realizations $E(q, G)$ and $B(q, G)$. Note that there is a continuous bijection $\overline{B}(q, G) \rightarrow B(q, G)/G^{ad}$ which is a homeomorphism if for example G is a compact Lie group.

Another basic variation of the constructions in this section can be introduced by using the free pro- p -group P_n on n generators instead of the free group F_n . Note that the p -adic integers $\mathbb{Z}_p$ is given by $P_1 = \mathbb{Z}_p$, and

$$\text{Hom}(\mathbb{Z}_p, G) = \varinjlim_m \text{Hom}(\mathbb{Z}/p^m, G),$$

can be identified with the elements in G of order a power of p .

The group P_n is not abelian for $n > 1$. Furthermore,

(1)

$$\text{Hom}(P_n/\Gamma^2(P_n), G) = \varinjlim_m \text{Hom}((\mathbb{Z}/p^m)^n, G)$$

is the set of ordered, commuting n -tuples of elements in G which all have finite order given by a power of p ,

- (2) $\text{Hom}(P_n, G)$ is the set of ordered n -tuples of elements in G which all have finite order given by a power of p , and
- (3) the quotient $\text{Hom}(P_n, G)/G$ corresponds to G -conjugacy classes of these n -tuples.

One can construct the associated simplicial spaces as limits of simplicial spaces, with realizations denoted by $E(q, G)_p$ and $B(q, G)_p$. The spaces associated to quotients by the conjugation action are also well-defined, and they are denoted by $\overline{E}(q, G)_p$ and $\overline{B}(q, G)_p$ respectively. It seems worthwhile to point out that the free group can also be replaced by its pro-finite completion $\widehat{F}_n$ in the constructions described in this paper, which leads to further interesting spaces.

Note that the automorphism group of the free group F_n will act naturally on the spaces $\text{Hom}(F_n/\Gamma^q, G)$ and thus on their homology, providing interesting representations. Similarly there are actions of $GL(n, \mathbb{Z})$ on the homology of the spaces $\text{Hom}(\mathbb{Z}^n, G)$. If instead the pro-finite completion $\widehat{F}_n$ is used, this leads to a homology representation of the automorphism group of $\widehat{F}_n$. Recall that faithful representations of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ in the automorphism group of the profinite completion $\widehat{F}_n$ for $n \geq 2$ have been constructed (see [6] and [11]). Thus the space $\text{Hom}(\widehat{F}_n/\Gamma^q, G)$ admits a natural action of both $\text{Aut}(\widehat{F}_n)$ as well as $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. In case G is a finite group, there are associated permutation representations of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ which will be considered in a sequel.

3. $B(q, G)$ AS A COLIMIT AND HOMOTOPY COLIMIT

This section deals primarily with the case when G is a finite group.

Definition 3.1. For a finite group G , let

$$\mathcal{N}_q(G) = \{A \subset G \text{ subgroup} \mid \Gamma^q(A) = \{1\}\}$$

The elements of $\mathcal{N}_q(G)$ are precisely all the subgroups of nilpotence class less than q . Note that this is a partially ordered set under inclusion, furthermore it is closed under conjugation by elements in G and under the process of taking subgroups (indeed if $H \subset K$ is a subgroup, then $\Gamma^i(H) \subset \Gamma^i(K)$ for all $i \geq 1$). Such a collection is often referred to as a *family* of subgroups in G . A group can be defined associated to these subgroups and the inclusions between them.

Definition 3.2. If G is a finite group, let $G(q) = \text{colim}_{A \in \mathcal{N}_q(G)} A$.

$\mathcal{N}_2(G)$ is the family of abelian subgroups in G , hence $G(2)$ is simply the colimit of all abelian subgroups in G . The sequence of groups $G(2), G(3), G(4), \dots$ captures information

relevant to the subgroups of G of increasing nilpotence class. Note that $G(\infty) = G$. Their classifying spaces will play a role in the analysis of the spaces $B(q, G)$.

Theorem 3.3. *Let G be a finite group, then for any $q \geq 2$, $B(q, G)$ can be constructed as the pushout of the realization of the diagram of simplicial spaces obtained by applying the functor $B_*(q, -)$ to $\mathcal{N}_q(G)$, yielding the identifications*

$$B(q, G) = \operatorname{colim}_{A \in \mathcal{N}_q(G)} BA.$$

Proof. For any subgroup $H \subset G$, there are natural inclusions

$$\operatorname{Hom}(F_n/\Gamma^q, H) \subset \operatorname{Hom}(F_n/\Gamma^q, G).$$

Given any homomorphism $\psi : F_n \rightarrow G$ with $\psi(\Gamma^q) = 1$, it follows that $K = \operatorname{Im} \psi$ belongs to $\mathcal{N}_q(G)$, and that $\psi \in \operatorname{Hom}(F_n/\Gamma^q, \operatorname{Im} \psi)$. Now if $A \in \mathcal{N}_q(G)$, then $\operatorname{Hom}(F_n/\Gamma^q, A) = \operatorname{Hom}(F_n, A)$ for all $n \geq 0$; combining these facts yields

$$B_*(q, G) = \bigcup_{A \in \mathcal{N}_q(G)} B_*(\infty, A).$$

This simplicial space is contained in $B_*(\infty, G)$, and the realizations BA are natural subspaces of BG . If $A \subset A'$ then $BA \subset BA'$; moreover $BA \cap BA' = B(A \cap A')$. Therefore the realization $|\bigcup_{A \in \mathcal{N}_q(G)} B_*(\infty, A)|$ of the union is precisely the space that results from identifying the realizations BA along their intersections i.e. the colimit, whence the result follows. $\square$

Colimits are not easy to handle; it is preferable to deal with a *homotopy colimit* whenever possible (see [22]). Under favorable circumstances they are weakly homotopy equivalent. This will occur if the diagram of spaces is a *free diagram* (see [9]); for a union this will hold if all the intersections of spaces in the diagram are used in the construction. This is automatically verified for the diagrams under consideration here, because of the nice properties of the functor $A \mapsto BA$ which have been outlined above. Thus there is a weak equivalence $B(q, G) \simeq \operatorname{hocolim}_{A \in \mathcal{N}_q(G)} BA$. Now the identity element in G makes all of the spaces in these diagrams *pointed spaces* (i.e. the classifying space $B\{1\}$ is the natural basepoint $*$). From this it follows that the fundamental group of $B(q, G)$ can be computed as a colimit:

$$\pi_1(B(q, G), *) \cong \operatorname{colim}_{A \in \mathcal{N}_q(G)} A \cong G(q).$$

This can be expressed more succinctly as follows. Let $\{M_i \mid i \in I\}$ denote a collection of maximal subgroups in $\mathcal{N}_q(G)$. Then

$$\pi_1(B(q, G), *) \cong *_{i \in I} M_i / (r_{ij})$$

where the r_{ij} are the relations coming from the inclusions $M_i \cap M_j \subset M_i$.

Having established that $B(q, G)$ is a homotopy colimit, there are some well-known methods from homotopy theory that can be used to study its homotopy type.

Proposition 3.4. *There is a natural fibration $B(q, G) \rightarrow BG(q)$ with fiber a simply-connected finite dimensional complex K_q .*

Proof. If $A \in \mathcal{N}_q(G)$, then there is a natural inclusion $A \subset G(q)$. This gives rise to a fibration $G(q)/A \rightarrow BA \rightarrow BG(q)$, where the fiber is a discrete coset space. Taking homotopy colimits and applying the results in [16] (see [9], page 180) yields a fibration

$$\operatorname{hocolim}_{A \in \mathcal{N}_q(G)} G(q)/A \rightarrow \operatorname{hocolim}_{A \in \mathcal{N}_q(G)} BA \rightarrow BG(q)$$

and as the fiber K_q is a homotopy colimit of a finite collection of discrete spaces it is necessarily finite dimensional. On the other hand $BG(q)$ is a $K(\pi, 1)$ and the map $B(q, G) \rightarrow BG(q)$ induces an isomorphism on fundamental groups, whence the result follows. $\square$

A basic question is that of determining under what conditions the fibre K_q is contractible.

Question: *If G is a finite group, are the spaces $B(q, G)$ Eilenberg–Mac Lane spaces of type $K(G(q), 1)$?*

In the following section examples will be given where this question admits an affirmative answer.

For any $q \geq 2$, the group $G(q)$ admits a natural surjection onto G , as every element of G is contained in an abelian subgroup, and these groups all belong to $\mathcal{N}_q(G)$ for $q \geq 2$. i.e. there is a group extension

$$1 \rightarrow T(q) \rightarrow G(q) \rightarrow G \rightarrow 1.$$

The kernel is torsion-free by construction, as every finite group in $G(q)$ necessarily embeds in G . The group $T(q)$ acts freely on the coset spaces $G(q)/A$, with quotient precisely G/A . Applying homotopy colimits, the fibrations

$$G(q)/A \rightarrow G/A \rightarrow BT(q)$$

give rise to a fibration

$$K_q \rightarrow \operatorname{hocolim}_{A \in \mathcal{N}_q(G)} G/A \rightarrow BT(q).$$

On the other hand the fibrations

$$G/A \rightarrow BA \rightarrow BG$$

give rise to a fibration

$$\operatorname{hocolim}_{A \in \mathcal{N}_q(G)} G/A \rightarrow B(q, G) \rightarrow BG.$$

Comparing this to the natural fibration $E(q, G) \rightarrow B(q, G) \rightarrow BG$ it can be seen that there is an equivalence

$$E(q, G) \simeq \operatorname{hocolim}_{A \in \mathcal{N}_q(G)} G/A.$$

and that

$$\pi_1(E(2, G)) \cong T(q)$$

which is a torsion-free group. Note that the *hocolim* term is a finite complex. These spaces can be assembled into a commutative diagram of fibrations:

$$\begin{array}{ccccc} K_q & \xlongequal{\quad} & K_q & & \\ \downarrow & & \downarrow & & \\ E(q, G) & \longrightarrow & B(q, G) & \longrightarrow & BG \\ \downarrow & & \downarrow & & \parallel \\ BT(q) & \longrightarrow & BG(q) & \longrightarrow & BG \end{array}$$

K_q is contractible if and only if $B(q, G) \simeq BG(q)$; this condition is equivalent to $E(q, G) \simeq BT(q)$.

The complexes K_q admit a $G(q)$ action with finite stabilizers, which restricts to a free action of the subgroup $T(q)$. The orbit space $K_q/T(q)$ is a finite complex with a natural G -action, and it is homotopy equivalent to $E(q, G)$. Now taking the quotient by $G(q)$ yields the equivalence

$$K_q/G(q) \simeq \operatorname{hocolim}_{A \in \mathcal{N}_q(G)} \{*\} \simeq |\mathcal{N}_q(G)| \simeq \{*\}$$

due to the fact that the nerve of the category is contractible, as it has a minimal element (the trivial subgroup $\{1\}$). On the other hand, there is an identification

$$B(q, G) \simeq EG(q) \times_{G(q)} K_q.$$

Note that the projection map $B(q, G) \rightarrow |\mathcal{N}_q(G)|$ is a rational equivalence (as the isotropy is finite) and hence it follows that $H^i(B(q, G); \mathbb{Q}) \cong 0$ for $i > 0$.

Remark. The spaces $B(q, G, p)$ can also be analyzed using the methods discussed in this section. In particular it is not hard to show that

$$B(q, G, p) = \operatorname{colim}_{K \in \mathcal{S}_p(G)} B(q, K, p)$$

where $\mathfrak{S}_p(G)$ is the poset of p -subgroups in G . This in turn can be further decomposed using the the classifying spaces of p -subgroups which are of p -nilpotence class less than q .

4. THE HOMOLOGY OF $B(q, G)$ WHEN G IS FINITE

The spaces $E(q, G)$ and $B(q, G)$ are constructed in a natural way, and seem to contain interesting information about the group G . For example the homology groups $H_*(E(q, G); \mathbb{Z})$ are natural $\mathbb{Z}G$ -modules and exploring their properties would seem like a worthwhile goal. This can be done explicitly in the case when G is a finite group.

Recall that if K_* is a simplicial set then one can define $\mathbb{Z}K_*$ as the free abelian group on the simplices of K_* . The face maps of K_* define boundary maps by the equation

$$\partial_n = \sum_{i=0}^n (-1)^i d_i : \mathbb{Z}K_n \rightarrow \mathbb{Z}K_{n-1}$$

and so $\mathbb{Z}K_*$ becomes a chain complex. The crux of this construction are the isomorphisms

$$H_*(\mathbb{Z}K_*) \cong H_*(\text{Sing}(|K_*|)) \cong H_*(|K_*|)$$

where $\text{Sing}(|K_*|)$ is the singular complex of $|K_*|$. Consider $\mathbb{Z}B_*(q, G)$, which can be thought of as a subcomplex of $\mathbb{Z}B_*(\infty, G)$; in fact

$$\mathbb{Z}B_*(q, G) = \bigcup_{H \in \mathcal{N}_q(G)} \mathbb{Z}B_*(\infty, H)$$

Note that $\mathbb{Z}B_0(q, G) = \mathbb{Z}$ and $\mathbb{Z}B_1(q, G) = \mathbb{Z}[G]$ (the free abelian group on G), so the chain complex looks like

$$\cdots \rightarrow \mathbb{Z}B_2(q, G) \xrightarrow{\partial_2} \mathbb{Z}[G] \xrightarrow{\partial_1} \mathbb{Z}$$

where $\partial_1 = 0$ and $\partial_2(x, y) = y - xy + x$. Define for each $q \geq 2$ a subgroup of $\mathbb{Z}[G]$ by

$$I_q(G) = \langle y - xy + x \mid \Gamma^q(\langle x, y \rangle) = 1, \text{ with } x, y \in G \rangle$$

Thus $I_2(G) \subseteq I_3(G) \subseteq \cdots \subseteq I_\infty(G)$, where $\Gamma^q = \{1\}$ when $q = \infty$. Then

$$H_1(B(q, G)) = \mathbb{Z}[G]/I_q(G)$$

and so there is a sequence of surjective maps

$$H_1(B(2, G)) \rightarrow H_1(B(3, G)) \rightarrow \cdots \rightarrow H_1(BG)$$

Corollary 4.1. *If G is a finite group and $q \geq 2$, then the $H_i(B(q, G); \mathbb{Z})$ are finite abelian groups for all $i > 0$, and their torsion only occurs at primes dividing the order of G .*

Proof. As observed previously, $\mathbb{Z}B_*(q, G)$ can be thought of as the union of the chain complexes generated by the maximal subgroups of class $< q$. Therefore a Mayer-Vietoris spectral sequence can be used to compute $H_*(B(q, G); \mathbb{Z})$, involving the homology of finite subgroups of G ; hence their reduced homology is annihilated by $|G|$. Thus the only torsion involved is at the primes dividing $|G|$. $\square$

Similarly one can consider $\mathbb{Z}E_*(q, G)$; in this case $\mathbb{Z}E_0(q, G) = \mathbb{Z}$, and the complex looks like

$$\cdots \rightarrow \mathbb{Z}E_2(q, G) \xrightarrow{\partial_2} \mathbb{Z}E_1(q, G) \xrightarrow{\partial_1} \mathbb{Z}[G]$$

where $\partial_1(a, x) = ax - a$ and $\partial_2(a, x, y) = (ax, y) - (a, xy) + (a, x)$. As the natural projection $E_*(q, G) \rightarrow B_*(q, G)$ is a simplicial map this yields a map of chain complexes and hence one of homology groups. Note that this map on H_1 takes the form

$$H_1(E(q, G)) \rightarrow H_1(BG)$$

$$(z, x) \mapsto x$$

at the chain level. As before, there is a sequence of surjective maps

$$H_1(E(2, G)) \rightarrow H_1(E(3, G)) \rightarrow \cdots \rightarrow H_1(EG).$$

The following result shows that the first homology group contains interesting information.

Proposition 4.2. *The Feit–Thompson Theorem is equivalent to the following result: for G of odd order, the homomorphism*

$$H_1(E(2, G); \mathbb{Z}) \rightarrow H_1(B(2, G); \mathbb{Z})$$

cannot be surjective. The same statement holds for $H_1(E(q, G); \mathbb{Z}) \rightarrow H_1(B(q, G); \mathbb{Z})$ for any finite q .

Proof. Consider the fibration

$$E(2, G) \rightarrow B(2, G) \rightarrow BG$$

and its associated 5-term exact sequence in homology; this yields an exact sequence of the form

$$H_2(B(2, G); \mathbb{Z}) \rightarrow H_2(BG; \mathbb{Z}) \rightarrow H_1(E(2, G); \mathbb{Z})_G \rightarrow H_1(B(2, G); \mathbb{Z}) \rightarrow G/[G, G] \rightarrow 0.$$

The Feit–Thompson Theorem says that every odd order group is solvable, which is equivalent to the condition that $G/[G, G] \neq 1$ for all G of odd order. This is precisely equivalent to the failure of surjectivity for the map $H_1(E(2, G); \mathbb{Z}) \rightarrow H_1(B(2, G); \mathbb{Z})$. $\square$

Note that $H_1(E(q, G); \mathbb{Z}) \cong T(q)/[T(q), T(q)]$. In the sequel examples will be given where the geometry of the spaces $E(q, G)$ will be explicitly determined (especially in the case $q = 2$) and the map on H_1 analyzed in some detail.

5. THE SPACES $B(q, G)$ FOR COMPACT, CONNECTED LIE GROUPS

In this section basic properties of the spaces $B(q, G)$ will be analyzed for compact, connected Lie groups. This will include a calculation of the rational cohomology of $B(2, G)$ as well as a splitting theorem for $\Omega(B(q, G))$.

For a compact, connected Lie group G , consider the map $\phi_n : G/T \times T^n \rightarrow \text{Hom}(\mathbb{Z}^n, G)$, given by

$$(g, t_1, \dots, t_n) \rightarrow (gt_1g^{-1}, \dots, gt_ng^{-1}).$$

Notice that the Weyl group $W(G) = N(T)/T$ acts freely on $G/T \times T^n$ by

$$(gT, t_1, \dots, t_n) \cdot w = (gwT, w^{-1}t_1w, \dots, w^{-1}t_nw)$$

and that the map ϕ_n is invariant under this action. This yields the following commutative diagram

$$\begin{array}{ccc} G/T \times T^n & \xrightarrow{\phi_n} & \text{Hom}(\mathbb{Z}^n, G) \\ \downarrow & \nearrow \overline{\phi_n} & \\ (G/T \times T^n)/W(G) & & \end{array}$$

Note that the vertical map is equivalent to the map $(G \times T^n)/T \rightarrow (G \times T^n)/N(T)$. Assume that the space $\text{Hom}(\mathbb{Z}^n, G)$ is path-connected, then as was shown in [5], ϕ_n is surjective and the fibers of the induced map

$$\overline{\phi_n} : (G \times T^n)/N(T) \rightarrow \text{Hom}(\mathbb{Z}^n, G)$$

are rationally acyclic. Hence applying the Vietoris-Begle Theorem it follows that $\overline{\phi_n}$ induces a rational homology equivalence and thus the map ϕ_n induces an isomorphism

$$H^*(\text{Hom}(\mathbb{Z}^n, G); \mathbb{Q}) \rightarrow H^*(G/T \times T^n; \mathbb{Q})^{W(G)}.$$

This information can be assembled to yield the following theorem.

Theorem 5.1. *Let G denote a compact, connected Lie group with maximal torus $T \subset G$. Assume that the spaces $\text{Hom}(\mathbb{Z}^n, G)$ are all path connected. There is an isomorphism*

$$H^*(B(2, G); \mathbb{Q}) \cong H^*(G/T \times BT; \mathbb{Q})^{W(G)}$$

which is compatible with the well-known isomorphism $H^(BG; \mathbb{Q}) \cong H^*(BT; \mathbb{Q})^{W(G)}$, where $W(G)$ denotes the Weyl group of T in G .*

Proof. Identify the space $G/T \times T^n$ with the product $G/T \times B_n(\infty, T)$, where G/T is seen as a constant simplicial space. Thus the maps ϕ_n define a simplicial map ϕ , the Weyl group $W(G)$ acts simplicially on $G/T \times B_*(\infty, T)$ and ϕ is invariant under this action. This yields the following commutative diagram of simplicial spaces

$$\begin{array}{ccc} G/T \times B_*(\infty, T) & \xrightarrow{\phi} & B_*(2, G) \\ \downarrow & \nearrow \bar{\phi} & \\ (G/T \times B_*(\infty, T))/W(G) & & \end{array}$$

The simplicial spaces under consideration are all proper (note that this was proved in [3] for $B_*(2, G)$) and so by a standard result (see [14]) the rational cohomology isomorphisms induced by the maps $\bar{\phi}_n$ induce an isomorphism between the rational cohomology of the geometric realizations, yielding the desired equivalence. The last assertion follows from the diagram

$$\begin{array}{ccc} G/T \times BT & \xrightarrow{\phi} & B(2, G) \\ \downarrow \text{proj} & & \downarrow \text{inc} \\ BT & \xrightarrow{\text{inc}} & BG \end{array}$$

which is commutative up to homotopy. $\square$

Remark. If the spaces $\text{Hom}(\mathbb{Z}^n, G)$ are not path-connected, an analogous result can be obtained using the generic component $\text{Hom}_0(\mathbb{Z}^n, G)$ i.e. the component of the trivial representation. This is indeed a simplicial subspace, as the face and degeneracy maps are continuous and preserve the trivial representation. Its realization gives rise to a subspace $B(2, G)_0 \subset B(2, G)$ for which the rational cohomology can be computed as above.

Example 5.2. In case $G = U(n)$, it was shown in [3] that the spaces of ordered commuting n -tuples are all path-connected. Hence the theorem applies here. Note that in this case the Weyl group is the symmetric group S_n which acts by permuting the entries in the diagonal maximal torus. Recall that

$$H^*(BT) \cong \mathbb{Q}[t_1, \dots, t_n],$$

a polynomial ring on n two dimensional generators. Thus the rational cohomology of $B(2, U(n))$ can be described as the ring of invariants

$$H^*(B(2, U(n)); \mathbb{Q}) \cong (H^*(U(n)/T) \otimes \mathbb{Q}[t_1, \dots, t_n])^{S_n}.$$

Note that the natural inclusion

$$B(2, U(n)) \longrightarrow BU(n)$$

has a stable section, as this is true for the map $BT \rightarrow BU(n)$, where $T \subset U(n)$ is a maximal torus [20]. Thus $BU(n)$ is a stable retract of $B(q, U(n))$ for all $q \geq 2$.

The spaces $Hom(\mathbb{Z}^n, G)$ admit an action of G by conjugation, and the orbit space $Rep(\mathbb{Z}^n, G)$ can be identified with the moduli space of flat G -bundles over the torus T^n , an object of considerable geometric interest.

The next result in this section is a product decomposition arising from looping the fibration

$$E(q, G) \rightarrow B(q, G) \rightarrow BG.$$

First recall that there is a map

$$\iota_G : \Sigma(G) \rightarrow BG$$

which is natural for morphisms of topological groups obtained by identifying the first filtration $F_1 BG$ as $\Sigma(G)$. In addition, the composite

$$G \xrightarrow{E} \Omega\Sigma(G) \xrightarrow{\Omega(\iota_G)} \Omega BG$$

is a homotopy equivalence where E is the standard Freudenthal suspension map and G is a Lie group.

The splitting result is stated next.

Theorem 5.3. *If G is a connected Lie group, then the associated looped fibration*

$$\Omega(E(q, G)) \rightarrow \Omega(B(q, G)) \rightarrow \Omega(BG)$$

has a cross-section (up to homotopy). The cross-section is induced by a map

$$\sigma(q, G) : G \rightarrow \Omega(B(q, G))$$

given by the composite

$$G \xrightarrow{E} \Omega(\Sigma(G)) \xrightarrow{\Omega(\iota(q, G))} \Omega(B(q, G))$$

where

$$\iota(q, G) : \Sigma(G) \rightarrow B(q, G)$$

is given by the canonical identification $\Sigma(G) = F_1 B(q, G)$ for $q \geq 2$, and which is natural for morphisms in G . Thus there is a homotopy equivalence

$$\theta(q, G) : G \times \Omega(E(q, G)) \rightarrow \Omega B(q, G).$$

Furthermore the cross-section $\sigma(q, G)$ and homotopy equivalence $\theta(q, G)$ are natural in the sense that if

$$f : G \rightarrow H$$

is a morphism of topological groups, then there are strictly commutative diagrams

$$\begin{array}{ccccc} G & \xrightarrow{\sigma(q, G)} & \Omega B(q, G) & \longrightarrow & \Omega BG \\ f \downarrow & & \downarrow \Omega B(q, f) & & \downarrow \Omega Bf \\ H & \xrightarrow{\sigma(q, H)} & \Omega B(q, H) & \longrightarrow & \Omega BH. \end{array}$$

and

$$\begin{array}{ccc} G \times \Omega(E(q, G)) & \xrightarrow{\theta(q, G)} & \Omega B(q, G) \\ f \times \Omega E(q, f) \downarrow & & \downarrow \Omega B(q, f) \\ H \times \Omega(E(q, H)) & \xrightarrow{\theta(q, H)} & \Omega B(q, H). \end{array}$$

Proof. Recall that there is a canonical identification $\Sigma(G) = F_1 B(q, G)$ for $q \geq 2$ natural for morphisms in G . Thus there is a natural factorization

$$\Sigma(G) \longrightarrow F_1 B(2, G) \longrightarrow \cdots \longrightarrow F_1(BG).$$

together with the associated composite arising by taking adjoints

$$G \xrightarrow{E} \Omega \Sigma(G) \longrightarrow \Omega F_1 B(2, G) \cdots \longrightarrow \Omega(F_1(BG)) \longrightarrow \Omega(BG).$$

So the following diagram strictly commutes:

$$\begin{array}{ccccccc} G & \xrightarrow{E} & \Omega \Sigma(G) & \longrightarrow & \Omega F_1 B(2, G) \cdots & \longrightarrow & \Omega(F_1(BG)) \longrightarrow \Omega(BG) \\ 1 \downarrow & & 1 \downarrow & & \downarrow & & \downarrow \\ G & \xrightarrow{E} & \Omega \Sigma(G) & \longrightarrow & \Omega B(2, G) \cdots & \longrightarrow & \Omega(B(q, G)) \longrightarrow \Omega(BG) \end{array}$$

The composite

$$G \xrightarrow{E} \Omega \Sigma(G) \longrightarrow \Omega B(2, G) \longrightarrow \cdots \longrightarrow \Omega(B(q, G))$$

is the adjoint $\iota(q, G) : \Sigma(G) \rightarrow B(q, G)$ and thus the induced map

$$G \xrightarrow{E} \Omega \Sigma(G) \longrightarrow \Omega(B(q, G)) \longrightarrow \Omega(BG)$$

is a homotopy equivalence.

The looped fibration $\Omega(E(q, G)) \rightarrow \Omega(B(q, G)) \rightarrow \Omega(BG)$ has a cross-section. The choice of homotopy equivalence $\theta(q, G) : G \times \Omega(E(q, G)) \rightarrow \Omega(B(q, G))$ is the composite

$$G \times \Omega(E(q, G)) \xrightarrow{\sigma(q, G) \times \Omega(p(q, G))} \Omega(B(q, G)) \times \Omega(B(q, G)) \xrightarrow{\text{multiply}} \Omega(B(q, G)).$$

Thus the total space $\Omega B(q, G)$ is homotopy equivalent to a product

$$G \times \Omega(E(q, G)).$$

To finish, it suffices to check naturality. Notice

(1) that the maps

$$\sigma(q, G) : G \rightarrow \Omega B(q, G)$$

are natural for morphisms in G ,

(2) the maps $\Omega(p(q, G)) : \Omega(E(q, G)) \rightarrow \Omega(B(q, G))$ are natural for morphisms in G , and

(3) the composite

$$G \times \Omega(E(q, G)) \xrightarrow{\sigma(q, G) \times \Omega(p(q, G))} \Omega(B(q, G)) \times \Omega(B(q, G)) \xrightarrow{\text{multiply}} \Omega(B(q, G))$$

is natural for morphisms in G .

The theorem follows. □

6. $B(2, G)$ FOR TRANSITIVELY COMMUTATIVE GROUPS

In this section the space $B(2, G)$ will be described for a very particular class of finite groups. The following example serves as a motivation.

Example 6.1. In the dihedral group $D_{2n} = \langle a, b | a^n = b^2 = 1, bab = a^{-1} \rangle$ the conjugacy classes are as follows

$$(a^r)^{D_{2n}} = \{a^r, a^{-r}\}, \quad (b)^{D_{2n}} = \{a^{2t}b | t \geq 0\}, \quad (ab)^{D_{2n}} = \{a^{2t+1}b | t \geq 0\}.$$

If n is odd then there is only one conjugacy class of elements of order two. If n is even, then the center $Z(D_{2n}) = \langle a^{n/2} \rangle$. The centralizers are as follows:

If $n = 2k + 1$, then

$$C(a^r) = \langle a \rangle, \quad C(b) = \langle b \rangle, \quad C(ab) = \langle ab \rangle.$$

If $n = 2k$, then

$$C(a^r) = \begin{cases} D_{2n} & \text{if } r = k \\ \langle a \rangle & \text{if } r \neq k \end{cases}$$

$$C(b) = \{1, b, a^k b, a^k\}, \quad C(ab) = \{1, ab, a^{k+1}b, a^k\}.$$

Let $(g_1, \dots, g_m) \in D_{2n}^m$ and without loss of generality assume that g_1 is not a central element. Then $(g_1, \dots, g_m) \in \text{Hom}(\mathbb{Z}^m, D_{2n})$ if and only if $g_1, g_2, \dots, g_m \in C(g_1)$, as the centralizers of noncentral elements are abelian.

Thus, if n is odd

$$\text{Hom}(\mathbb{Z}^m, D_{2n}) = \text{Hom}(\mathbb{Z}^m, C(a)) \cup \bigcup_{0 \leq i < n} \text{Hom}(\mathbb{Z}^m, C(a^i b)),$$

and these sets have $\text{Hom}(\mathbb{Z}^m, \{1\})$ as its pairwise intersection. Therefore

$$B(2, D_{2n}) = BC_n \vee \bigvee_n BC_2$$

A similar description holds when n is even.

The dihedral groups belong to a special class of groups. The following Lemma characterizes these groups.

Lemma 6.2. *Let G be a nonabelian group. The following are equivalent*

- a) *If $g \notin Z(G)$, then $C(g)$ is abelian.*
- b) *If $[g, h] = 1$, then $C(g) = C(h)$ whenever $g, h \notin Z(G)$.*
- c) *If $[g, h] = 1 = [h, k]$, then $[g, k] = 1$ whenever $h \notin Z(G)$.*
- d) *If $A, B \leq G$ and $Z(G) < C_G(A) \leq C_G(B) < G$, then $C_G(A) = C_G(B)$.*

Proof. (a) $\Rightarrow$ (b) $\Rightarrow$ (c) is immediate. (c) $\Rightarrow$ (d) Let $a \in A, g \in C_G(A) - Z(G)$, $b \in B - Z(G)$, and $h \in C_G(B)$. We have $[b, g] = 1 = [g, a]$, so $[b, a] = 1$. As $[h, b] = 1 = [b, a]$ it follows that $[h, a] = 1$, hence $C_G(B) \subseteq C_G(A)$ as wanted. (d) $\Rightarrow$ (a) Let $x \in G - Z(G)$ and $g, h \in C_G(x)$. Thus $Z(G) < C_G(x, g) \leq C_G(x) < G$, thus $C_G(x, g) = C_G(x)$ and hence $[g, h] = 1$. $\square$

Definition 6.3. A finite group satisfying any of the conditions of Lemma 6.2 is called a transitively commutative group, or simply a TC group.

Remark. TC-groups are classified, see [18]. Some examples include

- (1) Groups with an abelian normal subgroup of prime index, e.g. dihedral and generalized quaternion groups.
- (2) $SL(2, \mathbb{F}_{2^n})$, with $n \geq 2$.

(3) All nonabelian groups of order < 24 .

Let $a_1, \dots, a_k \in G - Z(G)$ be a set of representatives of their centralizers so that

$$G = \bigcup_{1 \leq i \leq k} C_G(a_i)$$

and no smaller number of centralizers covers G . Note that Lemma 6.2 shows that the groups defining this union do not depend on the choice of representatives. This number k is called the *number of centralizers that cover G* . Moreover, note that each $C(a_i)$ is a maximal abelian subgroup of G .

Recall the basic property: $C(x) \cap C(y) = C(\langle x, y \rangle)$. So it follows that if G is a TC group, then its lattice of abelian subgroups looks like

$$\begin{array}{ccc} C(a_1) & \cdots & C(a_k) \\ & \searrow \quad \swarrow & \\ & Z(G) & \end{array}$$

A consequence of this is the following

Theorem 6.4. *If G is a TC group, then $B(2, G) \simeq BG(2)$, where $G(2)$ is the amalgamated product of the maximal abelian subgroups of G along the center of G . In particular, $E(2, G)$ is a $K(\pi, 1)$ as well.*

Proof. The space $B(2, G)$ is obtained by taking the pushout over the diagram that results from applying the classifying space functor to the lattice of abelian subgroups of G . Thus, the space $B(2, G)$ can be constructed inductively over the lattice of abelian subgroups of G , that is, if $M_1, \dots, M_k$ are the maximal abelian subgroups of G , then

$$B(2, G) = BM_1 \cup_{BZ(G)} \operatorname{colim}_{A \in \mathcal{N}_2(G) - \{M_1\}} BA.$$

When $k = 2$ a Theorem of J. H. C. Whitehead (see [8], pages 50–51 for details) shows that $B(2, G)$ is a $K(M_1 *_Z M_2, 1)$. If $k > 2$, then

$$\operatorname{colim}_{A \in \mathcal{N}_2(G) \setminus \{M_1\}} BA$$

is a $K(\pi, 1)$ by inductive hypothesis, where π is the amalgamated product of $M_2, \dots, M_k$ along the center of G , and so by Whitehead's Theorem $B(2, G)$ is a $K(G(2), 1)$. $\square$

Subsequently properties of the group $G(2)$ will be obtained for G a finite TC group using the fact that it acts on a tree with finite isotropy. For now the focus is on determining the

algebraic structure of TC groups. The following lemma, for instance, leads to an explicit description of $B(2, G)$ for TC-groups with trivial center.

Lemma 6.5. *Let G be a TC group with trivial center.*

- a) *The Sylow subgroups of G are abelian and intersect trivially.*
- b) *$\{C_G(x) \mid x \notin Z(G)\}$ is the family of maximal abelian subgroups of G .*
- c) *The maximal abelian subgroups of G intersect trivially.*
- d) *If H is a maximal abelian subgroup and $P \in \text{Syl}_p(H)$, then $P \in \text{Syl}_p(G)$.*

Proof. For (a), let $P \in \text{Syl}_p(G)$ and $1 \neq x \in Z(P)$, thus $P \subseteq C_G(x)$ and so P is abelian. Let $P, Q \in \text{Syl}_p(G)$ so that there is $1 \neq x \in P \cap Q$. So $P, Q \subseteq C_G(x)$ and thus PQ is an abelian p -subgroup of G of order

$$|PQ| = \frac{|P||Q|}{|P \cap Q|} > |P|$$

which is a contradiction.

Parts (b) and (c) are immediate.

For (d), let $P \in \text{Syl}_p(H)$ and $Q \in \text{Syl}_p(G)$ so that $P \subseteq Q$. Let $1 \neq x \in P$, so $H, Q \subseteq C_G(x)$ and thus HQ is an abelian subgroup. It follows that $Q \subseteq H$ since H is a maximal abelian subgroup. The result follows. $\square$

Remark. Note that Part (b) of the previous lemma holds for any TC group.

With the notation of Theorem 6.4, the following result holds:

Corollary 6.6. *If G is a TC group with trivial center, then*

$$B(2, G) \simeq \bigvee_{1 \leq i \leq k} \left(\prod_{p \mid |C_G(a_i)|} BP \right)$$

where $P \in \text{Syl}_p(G)$.

Proof. $B_*(2, G)$ is the one-point union of the simplicial spaces $B_*(2, C_G(a_i))$, and by Lemma 6.5

$$C_G(a_i) \cong \prod_{p \mid |C_G(a_i)|} P$$

with $P \in \text{Syl}_p(G)$. The result follows. $\square$

Given two finite groups P and Q of coprime orders, the homomorphism $P * Q \rightarrow P \times Q$ induces a homology isomorphism (this is easily seen from the Künneth formula). Hence the two classifying spaces $B(P * Q)$ and $B(P \times Q)$ are stably homotopy equivalent. This observation can be used to prove

Proposition 6.7. *If G is a finite TC group with trivial center, then there is a stable homotopy equivalence*

$$B(2, G) \simeq \bigvee_{p||G|} \bigvee_{P \in \text{Syl}_p(G)} BP.$$

This result implies the following cohomology calculation:

Corollary 6.8. *If G is a TC group with trivial center, then for $i > 0$,*

$$H^i(B(2, G); \mathbb{Z}) = \bigoplus_{p||G|} \bigoplus_{n_p(G)} H^i(P; \mathbb{Z})$$

where $P \in \text{Syl}_p(G)$ and n_p is the number of Sylow p -subgroups of G .

Example 6.9. The alternating group A_5 is a TC group. Its p -Sylow subgroups are $\mathbb{Z}/2 \times \mathbb{Z}/2$, $\mathbb{Z}/3$ and $\mathbb{Z}/5$. Then there is a homotopy equivalence

$$B(2, A_5) \simeq \bigvee^5 B(\mathbb{Z}/2 \times \mathbb{Z}/2) \bigvee^{10} [\bigvee B\mathbb{Z}/3] \bigvee^6 [\bigvee B\mathbb{Z}/5].$$


Consider the structure of $E(2, G)$ for G a finite TC group. The main ingredient which can be applied is the theory of trees, as described in [19]. In this situation, the colimit group $G(2) = *_ {Z(G)} C_G(a_i)$ acts on a graph X by setting

$$\text{Edges}(X) = \coprod_{i=1}^k G(2)/Z(G)$$

$$\text{Vertices}(X) = G(2)/Z(G) \sqcup \left[\coprod_{1 \leq i \leq k} G(2)/C(a_i) \right]$$

and where the i^{th} copy of $G(2)/Z(G)$ is identified with the vertices $G(2)/Z(G) \sqcup G(2)/C(a_i)$ by using the identity and the natural maps $G(2)/Z(G) \rightarrow G(2)/C(a_i)$. It turns out that X is a tree (see [19] p. 38), with a simplicial action of $G(2)$ having as fundamental domain the

tree of groups given by



A corollary of this construction is a version of a theorem originally due to Kurosch (see for instance Corollary A.2 from [8] or [19] p. 56).

Lemma 6.10. *Let F be a subgroup of a free product of finite groups amalgamated along a common subgroup to all of the factors. If F intersects trivially with every conjugate of the factors, then F is free.*

In particular this can be applied to prove that the subgroup $T(q) = \pi_1(E(2, G))$ in $G(2)$ is a free group of finite rank.

Proposition 6.11. *If G is a TC group then the fundamental group of $E(2, G)$ is a free group of rank*

$$N_G = 1 - |G : Z(G)| + \left(\sum_{1 \leq i \leq k} |G : Z(G)| - |G : C_G(a_i)| \right)$$

and so $E(2, G) \simeq \bigvee_{N_G} S^1$.

Proof. Notice that the fundamental group of $E(2, G)$ is the kernel of the homomorphism between $G(2)$ to G induced by the natural inclusions, and no element in $T(2) = \pi_1(E(2, G))$ is conjugate to an element in a centralizer or else the image of this element is nontrivial in G . Hence by the lemma above, it must be a free group. To find the rank it suffices to calculate the Euler characteristic of the groups in the extension

$$1 \rightarrow T(2) \rightarrow G(2) \rightarrow G \rightarrow 1$$

On the one hand

$$\chi(G(2)) = \frac{\chi(T(2))}{|G|}$$

and so

$$\text{rank}(T(2)) = 1 - \chi(T(2)) = 1 - \chi(G(2))|G|.$$

On the other hand, since $G(2)$ is the amalgamated product of the centralizers covering G along $Z(G)$, it follows by an inductive argument that

$$\chi(G(2)) = \frac{1}{|Z(G)|} + \sum_{i=1}^k \left[\frac{1}{|C_G(a_i)|} - \frac{1}{|Z(G)|} \right]$$

$$\text{rank } T(2) = 1 - \chi(G(2))|G| = 1 - |G| \left(\frac{1}{|Z(G)|} + \sum_{1 \leq i \leq k} \left[\frac{1}{|C_G(a_i)|} - \frac{1}{|Z(G)|} \right] \right)$$

and the result follows. $\square$

Note that $Y = X/T(2)$ is a finite graph with a G -action. Looking at the cellular chain complex of Y yields the following exact sequence of $\mathbb{Z}G$ -modules

$$0 \rightarrow H_1(E(2, G)) \rightarrow \bigoplus_{i=1}^k \mathbb{Z}[G/Z(G)] \xrightarrow{\phi} \mathbb{Z}[G/Z(G)] \oplus \bigoplus_{i=1}^k \mathbb{Z}[G/C(a_i)] \rightarrow \mathbb{Z} \rightarrow 0 \quad (I)$$

where $\mathbb{Z}[G/H]$ denotes the usual permutation module with isotropy H , and the structure of $M(2, G) = H_1(E(2, G))$ as a G -module is the one defined by the extension

$$1 \rightarrow \pi_1(E(2, G)) \rightarrow G(2) \rightarrow G \rightarrow 1.$$

The map ϕ is described as follows; let $j_i : \mathbb{Z}[G/Z(G)] \rightarrow \mathbb{Z}[G/C(a_i)]$ be the natural projection induced by the inclusion of $Z(G)$ in $C(a_i)$ for $i = 1, \dots, k$. Then

$$\phi(v_1, v_2, \dots, v_k) = \left(\sum_{i=1}^k v_i, -j_1(v_1), \dots, -j_k(v_k) \right)$$

There are two long exact sequences associated to this exact sequence, obtained by applying G -hypercohomology to it (see [4], Chapter V). The first one is:

$$\dots \rightarrow H_i(G; \mathbb{Z}) \rightarrow H_{i-2}(G, M(2, G)) \rightarrow H_{i-1}(B(2, G); \mathbb{Z}) \rightarrow H_{i-1}(G; \mathbb{Z}) \rightarrow \dots \quad (II)$$

Note that this sequence terminates with

$$H_2(G) \rightarrow H_0(G, M(2, G)) \rightarrow H_1(B(2, G)) \rightarrow H_1(G) \rightarrow 0$$

The other one is of the form

$$\rightarrow \dots \rightarrow H_{q+1}(B(2, G)) \rightarrow \bigoplus_{i=1}^k H_q(Z(G)) \xrightarrow{\phi_*} H_q(Z(G)) \oplus \left[\bigoplus_{i=1}^k H_q(C(a_i)) \right] \rightarrow H_q(B(2, G)) \rightarrow \dots \quad (III)$$

From the definition of ϕ it is direct to verify that

$$\phi_*(x_1, \dots, x_k) = \left(\sum_{i=1}^k x_i, -j_{1*}(x_1), \dots, -j_{k*}(x_k) \right).$$

If A and B are finite abelian groups and $A \subset B$, then $H_*(A) \subset H_*(B)$, whence it follows that ϕ_* is injective, and so for each $q \geq 0$, $H_q(B(2, G))$ can be described via the short exact sequence

$$0 \rightarrow \bigoplus_{i=1}^k H_q(Z(G)) \xrightarrow{\phi_*} H_q(Z(G)) \oplus \bigoplus_{i=1}^k H_q(C(a_i)) \rightarrow H_q(B(2, G)) \rightarrow 0$$

In particular this yields the isomorphism

$$H_1(B(2, G)) \cong \{(Z(G) \oplus [\bigoplus_{i=1}^k C(a_i)])\} / \left(\phi_* \left(\bigoplus_{i=1}^k Z(G) \right) \right)$$

Now consider the special case when G is a TC group with a trivial center. In this situation the chain group $C_1(Y)$ is a free $\mathbb{Z}G$ -module, and the p -Sylow subgroups of the centralizers $C_G(a_i)$ are p -Sylow subgroups for G .

Lemma 6.12. *Let M denote a permutation $\mathbb{Z}G$ -module of finite rank and $\epsilon : M \rightarrow \mathbb{Z}$ the usual augmentation map onto the trivial module. Then the map ϵ splits over $\mathbb{Z}G$ if and only if $|G|$ divides the least common multiple of the orders of the isotropy subgroups.*

Proof. We can write $M = \bigoplus_{i=1}^k \mathbb{Z}[G/H_i]$ where the $H_i \subset G$ are subgroups. If the map ϵ splits, then $\mathbb{Z}/|G|$ is a direct summand in $\widehat{H}^0(G, M) \cong \bigoplus_{i=1}^k \mathbb{Z}/|H_i|$ hence $|G|$ divides the l.c.m. of the orders of the subgroups $H_1, \dots, H_k$. Conversely, suppose that $|G|$ divides this l.c.m. This means that it is possible to find a summand $\mathbb{Z}/|G| \subset \bigoplus_{i=1}^k \mathbb{Z}/|H_i|$ which maps bijectively onto $\widehat{H}^0(G; \mathbb{Z})$ under the map induced in cohomology by ϵ . Using the main result in [1], there exists a trivial submodule of rank one $T \subset M$ which represents this class of highest exponent and splits off as a direct summand of M ; this defines the desired splitting. $\square$

Consider the case when G is a TC group with trivial center. The kernel of the augmentation map $\epsilon : C_0(Y) \rightarrow \mathbb{Z}$ is isomorphic to the quotient module $C_1(Y)/M(2, G)$, where $C_1(Y)$ is a free $\mathbb{Z}G$ module. Applying the previous result yields a G -splitting:

$$C_0(Y) \cong \mathbb{Z} \oplus [C_1(Y)/M(2, G)]$$

For the statement of the following corollary recall that given M a $\mathbb{Z}G$ -module, a finitely generated projective module F , and a surjection $f : F \rightarrow M$, then the kernel of f is uniquely defined up to projective summands, and is denoted $\Omega^1(M)$ (see [4], Chapter II). It is referred to as a dimension-shift of M . Similarly if $N \rightarrow F'$ is a monomorphism and F'

is projective, then the cokernel is uniquely defined up to projective summands and denoted $\Omega^{-1}(N)$.

Corollary 6.13. *If G is a TC group with trivial center, then $M(2, G)$ is a $\mathbb{Z}$ -torsion free module which up to projective factors is isomorphic to $\Omega^1(I)$, where I denotes the kernel of the augmentation map $C_0(Y) \rightarrow \mathbb{Z}$. Furthermore there is a splitting of $\mathbb{Z}G$ -modules*

$$C_0(Y) \cong \mathbb{Z} \oplus [C_1(Y)/M(2, G)]$$

where up to projective summands $[C_1(Y)/M(2, G)] \cong \Omega^{-1}(M(2, G))$. Thus for $j \geq 1$

$$H_j(G, M(2, G)) \oplus H_{j+1}(G; \mathbb{Z}) \cong \bigoplus_{i=1}^k H_{j+1}(C(a_i), \mathbb{Z}) \cong H_{j+1}(B(2, G), \mathbb{Z})$$

Note that the homological statement in the corollary above is simply that the long exact sequence (II) splits. In more geometric terms, there is a cofiber sequence

$$B(2, G) \rightarrow BG \rightarrow W(2, G)$$

for which there is a stable splitting

$$\Sigma BG \vee W(2, G) \simeq \bigvee_{1 \leq i \leq k} \Sigma BC(a_i) \simeq \Sigma B(2, G)$$

where the homology of $W(2, G)$ is determined from the terms $H_*(G, M(2, G))$.

Example 6.14. Let $G = A_5$, the alternating group on five letters. This is a TC group with trivial center. The maximal abelian subgroups are the p -syllow subgroups, which are just $\mathbb{Z}/3$ (10 copies), $\mathbb{Z}/5$ (6 copies) and $\mathbb{Z}/2 \times \mathbb{Z}/2$ (5 copies). In this case $M(2, A_5)$ is a module of rank equal to 854 and

$$H_i(B(2, A_5)) \cong [H_i(\mathbb{Z}/2 \times \mathbb{Z}/2)]^5 \oplus [H_i(\mathbb{Z}/3)]^{10} \oplus [H_i(\mathbb{Z}/5)]^6 \cong H_{i-1}(A_5, M(2, A_5)) \oplus H_i(A_5).$$

Example 6.15. Let $G = SL_2(\mathbb{F}_8)$, this is a group of order $504 = 2^3 \cdot 3^2 \cdot 7$, with trivial centre and with p -syllow subgroups $(\mathbb{Z}/2)^3$, $\mathbb{Z}/9$ and $\mathbb{Z}/7$. In this case

$$H_i(B(2, SL_2(\mathbb{F}_8))) \cong [H_i((\mathbb{Z}/2)^3)]^9 \oplus [H_i(\mathbb{Z}/9)]^{28} \oplus [H_i(\mathbb{Z}/7)]^{36}$$

Example 6.16. Let $G = Q_8$, the quaternion group of order eight. In this case the centre is $\mathbb{Z}/2$, and there are three maximal abelian subgroups (each of order four) intersecting along this central subgroup. In this case $B(2, Q_8) \cong B(*_{\mathbb{Z}/2}^3 \mathbb{Z}/4)$, the amalgamation of the three

groups along the common $\mathbb{Z}/2$. The module $M(2, Q_8)$ is of rank equal to three. It fits into an exact sequence

$$0 \rightarrow M(2, Q_8) \rightarrow \mathbb{Z}[Q_8/\mathbb{Z}/2]^3 \rightarrow \mathbb{Z}[Q_8/\mathbb{Z}/2] \oplus \bigoplus_{i=1}^3 \mathbb{Z}[Q_8/C(a_i)] \rightarrow \mathbb{Z} \rightarrow 0$$

and $H_i(B(2, Q_8))$ can be computed from the sequence

$$0 \rightarrow [H_i(\mathbb{Z}/2)]^3 \rightarrow H_i(\mathbb{Z}/2) \oplus [H_i(\mathbb{Z}/4)]^3 \rightarrow H_i(B(2, Q_8)) \rightarrow 0.$$

Note that $H_i(B(2, Q_8)) = 0$ for i even. For i odd the map can be described as follows: let v_1, v_2, v_3 be the natural basis for $(\mathbb{Z}/2)^3$ and v, e_1, e_2, e_3 the natural generating set for $\mathbb{Z}/2 \oplus (\mathbb{Z}/4)^3$. Then the map between them is given by

$$v_1 \mapsto v + 2e_1, v_2 \mapsto v + 2e_2, v_3 \mapsto v + 2e_3$$

which has cokernel equal to $H_1(B(2, Q_8)) \cong \mathbb{Z}/4 \oplus \mathbb{Z}/2 \oplus \mathbb{Z}/2$. This can be extended to show

$$H_i(B(2, Q_8); \mathbb{Z}) \cong \begin{cases} \mathbb{Z} & \text{if } i = 0 \\ \mathbb{Z}/4 \oplus \mathbb{Z}/2 \oplus \mathbb{Z}/2 & \text{if } i > 0 \text{ odd} \\ 0 & \text{if } i > 0 \text{ even} \end{cases}$$

Note that the sequence (I) also allows us to compute the character of the representation

$$G \rightarrow \text{Aut}(H_1(E(2, G) \otimes \mathbb{C}) = GL(N_G, \mathbb{C}),$$

which will be denoted by $\mathcal{X}_{E(2, G)}$. Indeed, tensoring throughout with $\mathbb{C}$ yields a split sequence of modules from which information about this character can be easily obtained.

Example 6.17. For the dihedral group D_{2n}

$$\text{rank}(\pi_1(E(2, D_{2n}))) = \begin{cases} n^2 - 1 & n \text{ odd}, \\ k^2 - 1 & n = 2k. \end{cases}$$

Using the character table of D_{2n} it follows that if n is odd, then

$$\mathcal{X}_{E(2, G)} = (n - 1)X_2 + nY_1 + \cdots + nY_k,$$

where X_2 is the nontrivial 1-dimensional character of D_{2n} , and the Y_i 's are the 2-dimensional irreducible characters of D_{2n} .

The exact sequence of G -modules (I) yields

Proposition 6.18. *If G is TC group then the kernel of the character $\mathcal{X}_{E(2,G)}$ is precisely the center of G . Thus*

$$G/Z(G) \rightarrow \text{Aut}(H_1(E(2, G)))$$

is a faithful representation.

Note that this result shows that the representation $G \rightarrow \text{Out}(\pi_1(E(2, G)))$ is faithful when G is a TC group with trivial center; moreover, the representation

$$\pi_1(B(2, G)) \rightarrow \text{Aut}(\pi_1(E(2, G)))$$

is faithful as well.

In a more general setting it seems quite difficult to determine the homotopy type of $B(2, G)$. However in some instances $B(q, G)$ for $q > 2$ can be easier to describe.

Definition 6.19. Let G denote a finite group and let $\mathcal{P}_q(G)$ be the category with objects the set

$$\{M_\alpha, M_\alpha \cap M_\beta \mid M_\alpha, M_\beta \in \mathcal{M}_q(G)\}$$

where $\mathcal{M}_q(G)$ denotes the set of maximal subgroups in G of class $< q$, and the morphisms are the set of inclusions of the form $M_\alpha \cap M_\beta \rightarrow M_\alpha$.

This category is 1-dimensional, as there are no compositions. This can be identified with a *graph of groups*, that is, an oriented graph with a group at each vertex and a homomorphism between the two vertices of each edge according to orientation. Note that in this situation the graph is always connected. Now if G is a TC group, then the graph defined by $\mathcal{P}_2(G)$ is a tree, with arrows all pointing out from a single vertex which corresponds to the center of G . In fact the following general result holds:

Theorem 6.20. *Let $q \geq 2$ and G a finite group such that $\mathcal{P}_q(G)$ is a tree. Then the space $B(q, G)$ is aspherical and there is a natural homotopy equivalence*

$$B(q, G) \simeq BG(q).$$

Proof. Using the graph of groups $\mathcal{P}_q(G)$ a space $\mathbb{BP}(\mathcal{G})$ can be constructed by inserting a copy of the classifying space of each group at each vertex of the graph and by filling in a mapping cylinder for each map induced on classifying spaces on each edge (in this case arising from subgroup inclusions). The two ends of the mapping cylinder are identified with the classifying spaces on the respective vertices. This is a special case of a general construction for graphs of groups; in fact by [10], Theorem 1B.11 it follows that $\mathbb{BP}_q(G)$ is aspherical (the relevant hypothesis is the injectivity of all the homomorphisms on edges).

By construction the maps between classifying spaces have been replaced by cofibrations. Given that $\mathcal{P}_q(G)$ is a tree, the space $\mathbb{B}\mathcal{P}_q(G)$ is homotopy equivalent to their colimit (by collapsing the mapping cylinders) and this is precisely $B(q, G)$. $\square$

It is interesting to note that in the situation above, the colimit group $G(q)$ can be readily understood using the theory of trees as in [19]; the special condition can be described as those which determine a *tree of groups*, and $G(q)$ is the corresponding inductive limit associated to the tree.

Example 6.21. The symmetric group Σ_4 is a minimal example (i.e. of smallest order) of a non-TC group. The diagram of abelian subgroups is quite intricate although it can be shown that in fact $B(2, \Sigma_4)$ is aspherical. On the other hand the maximal subgroups of class less than 3 in Σ_4 are isomorphic to either D_8 or $\mathbb{Z}/3$, and every pair of distinct copies of D_8 intersect along the subgroup

$$K = \{1, (12)(34), (13)(24), (23)(14)\} \cong \mathbb{Z}/2 \times \mathbb{Z}/2.$$

Furthermore, $\mathcal{N}_q(\Sigma_4) = \mathcal{N}_3(\Sigma_4)$ for all $q \geq 3$, hence it follows from the theorem above that

$$B(q, \Sigma_4) \simeq \bigvee_{A \in \mathcal{N}_q(\Sigma_4)}^4 B\mathbb{Z}/3 \vee B(*_K^3 D_8) \simeq B \operatorname{colim}_{A \in \mathcal{N}_q(\Sigma_4)} A$$

7. STABLE SPLITTINGS AND THE CARDINALITY OF $\operatorname{Hom}(F_n/\Gamma^q, G)$ FOR FINITE GROUPS

A stable splitting was given for the space of commuting n -tuples in [3]. Analogous stable splittings arise for $\operatorname{Hom}(F_n/\Gamma^q, G)$ from the fat wedge filtration of the product G^n where the base-point of G is 1_G .

Definition 7.1. As before, let $F_j G^n$ denote the subspace of G^n with at least j coordinates equal to 1_G and define subspaces of $\operatorname{Hom}(F_n/\Gamma^q, G)$ by the formula $S_n(j, q, G) = \operatorname{Hom}(F_n/\Gamma^q, G) \cap F_j G^n$. Let $S_n(q, G)$ denote $S_n(1, q, G)$. A Lie group G is said have *cofibrantly filtered elements* if the natural inclusions $I_j : S_n(j, q, G) \rightarrow S_n(j-1, q, G)$ are cofibrations for all n, q and j for which both spaces are non-empty.

It would seem that many Lie groups G should have cofibrantly filtered elements; it seems plausible to conjecture that this holds if G is a closed subgroup of $GL(n, \mathbb{C})$. Note that in [3] the weaker condition of having cofibrantly commuting elements (i.e. the special case $q = 2$) was indeed verified for these groups. The following result describes the stable structure of the spaces of homomorphisms $\operatorname{Hom}(F_n/\Gamma^q, G)$ in terms of more recognizable pieces.

Theorem 7.2. *If G has cofibrantly filtered elements, then there are homotopy equivalences*

$$\Sigma \text{Hom}(F_n/\Gamma^q, G) \rightarrow \bigvee_{1 \leq k \leq n} \Sigma \bigvee_{\binom{n}{k}} \text{Hom}(F_k/\Gamma^q, G)/S_k(q, G).$$

Looking at cohomology yields

Corollary 7.3. *If G has cofibrantly filtered elements, then there are isomorphisms of graded abelian groups*

$$H^*(\text{Hom}(F_k/\Gamma^q, G); \mathbb{Z}) \rightarrow H^*\left(\bigvee_{1 \leq k \leq n} \bigvee_{\binom{n}{k}} \text{Hom}(F_k/\Gamma^q, G)/S_k(q, G); \mathbb{Z}\right).$$

It turns out that the pieces in the decomposition above can in fact be identified (stably) in terms of the natural *filtration quotients* for the geometric realization of $B(q, G)$. This requires recalling the notion of a proper simplicial space.

Definition 7.4. A pair of spaces (X, A) is said to be a *strong NDR pair* provided that there are maps $u : X \rightarrow [0, 1]$ and a homotopy $h : X \times [0, 1] \rightarrow X$ such that (X, A) is an NDR pair, namely

- (1) $A = u^{-1}(0)$,
- (2) $h(x, 0) = x$ for all $x \in X$,
- (3) $h(a, t) \in A$ for all $(a, t) \in A \times [0, 1]$,
- (4) if $u(x) < 1$ then $h(x, 1) \in A$

and if $u(x) < 1$ then $u(h(x, t)) < 1$.

Definition 7.5. A simplicial space X_* is said to be *proper* if each pair $(X_n, S(X_n))$ is a strong NDR-pair for all n , where $S(X_n) = \cup s_i(X_{n-1}) \subset X_n$.

The following was proved as Lemma 11.3 in [14].

Proposition 7.6. *Assume that the simplicial space X_* is proper. Then the geometric realization $|X_*|$ is naturally filtered by $F_j|X_*|$ with induced homeomorphisms*

$$\Sigma^j(X_j/S(X_j)) \rightarrow F_j|X_*|/F_{j-1}|X_*|.$$

It follows from a direct modification of the arguments in [3] that if G has cofibrantly filtered elements, then the simplicial spaces $X_* = \text{Hom}(F_*/\Gamma^q, G)$ are proper, where (using our previous notation) $S(X_n) = S_n(1, q, G) = S_n(q, G)$. Thus this yields

Theorem 7.7. *If G has cofibrantly filtered elements then the natural filtration quotients $E_k^0(B(q, G)) = F_k B(q, G)/F_{k-1} B(q, G)$ of the geometric realization $B(q, G)$ are stably homotopy equivalent to the summands $\text{Hom}(F_k/\Gamma^q, G)/S_k(q, G)$.*

Remark. The results in this section are special cases of very general splittings for simplicial spaces X_* which are proper and simplicially NDR. Details of this appear in [2]; the main work is to verify these conditions for the simplicial spaces $B_*(q, G)$.

If G is a discrete group, the maps $I_j : S_n(j, q, G) \rightarrow S_n(j-1, q, G)$ of Definition 7.1 are all inclusions of pointed, discrete spaces, and are thus cofibrations. In particular the stable splittings in this section will hold for these groups.

Theorem 7.8. *If G is a discrete group, then G has cofibrantly filtered elements, and the natural filtration quotients $E_k^0(B(q, G)) = F_k B(q, G)/F_{k-1} B(q, G)$ of the geometric realization $B(q, G)$ are stably homotopy equivalent to the summands $\text{Hom}(F_k/\Gamma^q, G)/S_k(q, G)$. Thus there are stable homotopy equivalences*

$$\text{Hom}(F_n/\Gamma^q, G) \simeq \bigvee_{1 \leq k \leq n} \bigvee_{\binom{n}{k}} \Sigma^{-k}(E_k^0(B(q, G))).$$

Proof. There are homotopy equivalences

$$\Sigma \text{Hom}(F_n/\Gamma^q, G) \simeq \bigvee_{1 \leq k \leq n} \Sigma \bigvee_{\binom{n}{k}} \text{Hom}(F_k/\Gamma^q, G)/S_k(q, G).$$

Now Proposition 7.6 gives homotopy equivalences

$$\Sigma^k(\text{Hom}(F_k/\Gamma^q, G)/S_k(q, G)) \simeq E_k^0(B(q, G))$$

and thus stable homotopy equivalences

$$\text{Hom}(F_k/\Gamma^q, G)/S_k(q, G) \simeq \Sigma^{-k}(E_k^0(B(q, G))).$$

Combining these results completes the proof. $\square$

These decompositions will apply to the special case of a finite group with the discrete topology. Of course in this case the spaces of homomorphisms are finite sets, and the decompositions above can be interpreted as numerical formulae. The following definitions will be helpful for keeping track of the numbers which will appear in the formulae for the cardinality of these spaces of homomorphisms.

Definition 7.9. Let G denote a finite group. The integer $\lambda_n(q, G)$ is defined as the cardinality of $\text{Hom}(F_n/\Gamma^q, G)$, and the integer $\mu_k(q, G)$ is defined as the rank of $H_k(E_k^0(B(q, G); \mathbb{Z}))$.

An immediate consequence of Theorem 7.8 is

Corollary 7.10. *If G is a finite group, then*

$$\lambda_n(q, G) = 1 + \sum_{1 \leq k \leq n} \binom{n}{k} \mu_k(q, G).$$

The special case $q = 2$ is especially interesting as the formula above provides information on the cardinality of the set of commuting elements in a finite group. Below are some calculations for abelian and transitively commutative groups.

Example 7.11. Let A denote a finite abelian group, then $\text{Hom}(\mathbb{Z}^n, A) \cong A^n$. The preceding formula expresses this quantity in a particular form. Recall that $S_k(2, A)$ denotes the elements in A^n where at least one coordinate is equal to 1. Thus the cardinality of $\text{Hom}(\mathbb{Z}^k, A)/S_k(2, A)$ is given by $1 + (|A| - 1)^k$. Now $B(2, A) = BA$ and the rank of $H_k(E_k^0(B(2, A); \mathbb{Z}))$ is $(|A| - 1)^k$ (recall that the k -th homology group of a k -fold suspension of a finite set has rank one less than the cardinality of the set). Therefore $\mu_k(2, A) = (|A| - 1)^k$ and the formula becomes

$$|A|^n = 1 + \sum_{1 \leq k \leq n} \binom{n}{k} (|A| - 1)^k.$$

Example 7.12. Consider the case when G is a finite transitively commutative group with trivial center. Now the space $B_*(2, G)$ is the one-point union of the simplicial spaces $B_*C_G(a_i)$, $1 \leq i \leq N$ where each $C_G(a_i)$ is a maximal abelian subgroup and so in this case

$$\mu_k(2, G) = \sum_{1 \leq i \leq N} (|C_G(a_i)| - 1)^k$$

and

$$\lambda_n(2, G) = 1 + \sum_{1 \leq k \leq n} \binom{n}{k} \sum_{1 \leq i \leq N} (|C_G(a_i)| - 1)^k$$

If $G = A_5$, the alternating group, then there are three isomorphism classes of centralizers: $\mathbb{Z}/2 \times \mathbb{Z}/2$ (five copies), $\mathbb{Z}/3$ (ten copies) and $\mathbb{Z}/5$ (six copies). Using this yields

$$\mu_k(2, A_5) = 5 \cdot 3^k + 10 \cdot 2^k + 6 \cdot 4^k$$

and

$$\lambda_n(2, A_5) = 1 + \sum_{1 \leq k \leq n} \binom{n}{k} [5 \cdot 3^k + 10 \cdot 2^k + 6 \cdot 4^k]$$

On the other hand these formulae can be rearranged to express

$$\lambda_n(2, G) = 1 + \sum_{1 \leq i \leq N} [\lambda_n(2, C_G(a_i)) - 1]$$

which can also be deduced directly from the structure of the group.

REFERENCES

- [1] A. Adem, *Cohomological exponents of ZG-lattices*, J. Pure Appl. Algebra **58**(1989), no. 1, 1–5.
- [2] A. Adem, M. Bendersky, A. Bahri, F. R. Cohen and S. Gitler, *On stable decompositions for simplicial spaces*, preprint 2008.
- [3] A. Adem, F. Cohen, *Commuting Elements and Spaces of Homomorphisms*, Mathematische Annalen **338**, 587–626.
- [4] A. Adem, R.J. Milgram, **Cohomology of Finite Groups**, Springer-Verlag Grundlehren 309 (2004).
- [5] T. Baird, *Cohomology of the space of commuting n -tuples in a compact Lie group*, Algebr. Geom. Topol. **7** (2007), 737–754.
- [6] V. Belyi, *On Galois extensions of a maximal cyclotomic field*, Math. USSR Izv., **14**(1980), 247–256.
- [7] A. Borel, R. Friedman, J. Morgan, *Almost Commuting Elements in Compact Lie Groups*, Mem. Amer. Math. Soc. **157** (2002), no. 747, x+136 pp.
- [8] K. Brown, **Cohomology of Groups**, Springer-Verlag GTM 87 (1982).
- [9] E. Dror-Farjoun, **Cellular Spaces, Null Spaces and Homotopy Localization**, Springer LNM 1622 (1996).
- [10] A. Hatcher, **Algebraic Topology**, Cambridge University Press (2002).
- [11] Y. Ihara, *On the embedding of $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ in $\widehat{GT}$* , in: The Grothendieck Theory of Dessins d’Enfants, London Mathematical Society Lecture Notes Vol. 200, Cambridge University Press, 1994.
- [12] S. Mac Lane, *The Milgram bar construction as a tensor product of functors*, pp. 135–152, in Lecture Notes in Mathematics, Vol. 168 Springer, Berlin.
- [13] J. P. May, *The Geometry of Iterated Loop Spaces*, Lect. Notes in Math., **271**, Springer-Verlag, Berlin, 1972.
- [14] J. P. May *E_∞ spaces, group completions, and permutative categories*, New developments in topology (Proc. Sympos. Algebraic Topology, Oxford, 1972), pp. 61–93. London Math. Soc. Lecture Note Ser., No. 11, Cambridge Univ. Press, London, 1974.
- [15] J. McCleary, **A user’s guide to spectral sequences** Cambridge Studies in Advanced Mathematics, 58. Cambridge University Press, Cambridge, 2001.
- [16] V. Puppe, *A remark on homotopy fibrations*, Manuscripta Math. **12** (1974) 113–120.
- [17] D. Quillen, B.B. Venkov, *Cohomology of finite groups and elementary abelian subgroups*, Topology **11** (1972), 317–318.
- [18] R. Schmidt, *Subgroup lattices of groups*, de Gruyter Expositions in Mathematics 14, Berlin, 1994.

- [19] J.-P. Serre, **Trees**, Springer-Verlag, New York, 1980.
- [20] V. Snaith, **Algebraic cobordism and K-theory**, Mem. Am. Math. Soc. 221 (1979).
- [21] E. Torres-Giese, **Spaces of Homomorphisms and Group Cohomology**, Ph.D. Thesis, University of British Columbia (2007).
- [22] V. Welker, G. Ziegler, R. Živaljević, *Homotopy colimits-comparison lemmas for combinatorial applications*, J. Reine Angew. Math., 509(1999), 117-149.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER BC V6T 1Z2,
CANADA

E-mail address: `adem@math.ubc.ca`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ROCHESTER, ROCHESTER NY 14627, USA

E-mail address: `cohf@math.rochester.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR MI 48109, USA

E-mail address: `etorresg@umich.edu`